Principles of plasticity in the developing brain



Bryan Kolb Canadian Centre for Behavioural Neuroscience University of Lethbridge, Canada

Take home Message

Preconceptual, prenatal, early postnatal, and adolescent events alter brain development and brain function for a lifetime.

Understanding the nature of these events is important for designing interventions, education, and public policy.

The Questions

How does the brain develop?

How is the development of the brain affected by prenatal and postnatal experiences?

What might the mechanism(s) of the effects be?

How do early experiences influence later educational/social/health outcomes in children?


Developing the cerebral hemispheres



86 billion neurons 10¹⁴ connections

An engineering marvel...

Your brain is sculpted by a lifetime of experiences, especially in the first few years of life and during adolescence.



The Logic of Lab Animal Studies

Show functional changes with a wide range of behavioural measures taken at different ages

Infer circuit changes from Golgi stains

Infer mechanisms from epigenetics



What events in brain development are altered by postnatal experience?

- 1. Neurogenesis in infant/toddler period.
- 2. Synapse formation in infant/toddler period.
- 3. Pruning of cells and connections, especially in early adolescence.

Factors influencing brain development


- 1. sensory & motor experience
- 2. language & music
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- 10. perinatal injury
- **11. immune system**



Both gestational and infant tactile stimulation alter brain and behavioural development.

These effects can be seen in the adult brain and behaviour.

Skilled Reaching Success in Adulthood After Infant TS





Infant tactile stimulation improves adult motor & cognitive function.

This is correlated with increased synaptic space across the cortex.

How does this work?



FGF-2


Events that alter FGF-2 therefore should influence brain development. They do.

Factors influencing brain development

- 1. sensory & motor experience
- 2. language & music
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- 10. perinatal injury
- 11. immune system

Early Language Exposure Changes the Brain

My main point here is that language and cognitive activity (including music) change the trajectory of developing brain regions, and later behaviour.



New Zealand Education Study

Gave reading and mathematics tests to the same children at age 5 & 14.

Result: There was virtually no difference in relative standing – even with 8 years of schooling!

THUS: little change in school outcomes after age 5! Trajectories are set early...


Early Music Exposure Changes the Brain

Music training increases brain volume, improves school performance, increases IQ ~ 7 points

And, increases 'cognitive reserve' in aging...

Factors influencing brain development


- 1. sensory & motor experience
- 2. language & cognitive experience
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- **10.** perinatal injury
- **11. immune system**






By age 4, the low SES kids have a 30 million word exposure deficit...

It is not only vocabulary but also grammatical structure that is different... (Erica Hoff, 2003)



Poverty reduces OFC volume in a 25 year prospective study




Holz et al., Neurpsychopharmacology, 2015

Factors influencing brain development

- 1. sensory & motor experience
- 2. language & cognitive experience
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- 10. perinatal injury
- **11. immune system**


Adult Stress and PFC

d (A



Means what?

This recovers with time in young but not older animals...



Gestational Stress

Impaired motor, socio-emotional, & cognitive behaviour in adulthood.

Increased corticosterone levels in adulthood & increased corticosterone reactivity to even mild stressors...





Gestational Stress increases anxiety, especially in males


> **Gestational Stress feminizes** male play


Halliwell, Gibb & Kolb, in progress, 2016



spatial learning

Halliwell, Gibb & Kolb, in progress, 2016






Effects of Gestational Stress on Brain

- **1.** *Smaller* **brains in adulthood**
- 2. Abnormal connectivity

The Quebec Ice Storm of 1998

- One of the worst natural disasters in Canadian recorded history: power failure for up to 6 weeks in the coldest month of the year (~-20C).
- Many pregnant women at various stages of pregnancy and varying degrees of hardship.
- Children are being followed with behavior, MRIs, & epigenetics.




The Quebec Ice Storm of 1998

Behavior: cognitive (IQ), linguistic, motor, and play abnormalities measured at 2, 5.5, 8.5, and 11.5 years.

MRIs: still being analyzed


DP Laplante, S King and colleagues


1998 QC Ice Storm and methylation 13 years later



Hypere 1. Differentially methylated CGs responding to cognitive appraisal level. Heatmap showing methylation of the 500 most differentially methylated CGs (P < 0.003, FDR < 0.055) across all 34 individuals. Each column represents an individual and each row a single CG. A color

Gestational Stress hastens the onset of AD symptoms





Paternal Preconception Stress



30 min 2X per day 27 days prior to mating

Harker et al., Neuroscience, 2015



The effect of Paternal stress is similar to having a perinatal brain injury...

(e.g., Harker et al., (2015) Neuroscience, 303, 200-210.)

Paternal Stress and PFC


Main overall effect is reduced synaptic space

Biggest changes in OFC Smallest changes in HPC

Effects are larger in females... -different than gestational stress where M>F

Proposed to be transferred by noncoding RNAs in sperm...





This is different from adult stress in mPFC Harker et al., 2015; 2017


Stress Summary

Preconceptual, gestational, and postnatal stress all change behaviour, epigenetics, and neuronal morphology, although in different ways. This is very different from adult stress.

Developmental stress also reduces plasticity later in life.

Factors influencing brain development

- 1. sensory & motor experience
- 2. language & cognitive experience
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- 10. perinatal injury
- 11. immune system



All classes of drugs, including Rx, legal, and illicit drugs change PFC structure & functioning. Effects are especially large with prenatal exposure. (see review by Robinson & Kolb, *Neuropharmacol*, 2004)

All Psychoactive Drugs Leave a Footprint

Early exposure to drugs with chronic effects

Amphetamine Nicotine⁺ Caffeine⁺ Morphine Valproate⁺ Antidepressants ⁺ Methylphenidate Alcohol⁺ Antipsychotics⁺





Mother - 1 st generation Fetus - 2nd generation Reproductive cells - 3rd generation

Prenatal nicotine is associated with an increased response to nicotine and reduced cerebral plasticity in adulthood.

Fluoxetine (Prozac) is worse! E.G. Muhammad et al., *Synapse*, 2012

Factors influencing brain development

- 1. sensory & motor experience
- 2. language & cognitive experience
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- **10. perinatal injury**
- **11. immune system**


Foreign Adoption Studies

Neuroimage. 2015 January 15; 105: 112-119. doi:10.1016/j neuroimage.2014.10.020.

Duration of Early Adversity and Structural Brain Development in Post-Institutionalized Adolescents

Amanda S. Hodel, Ruskin H. Hunt, Raquel A. Cowell¹, Sara E. Van Den Heuvel, Megan R. Gunnar, and Kathleen M. Thomas Institute of Child Development, University of Minnesota, 51 East River Road, Minneapolis, MN, 55455, USA

Children adopted at mean age of 12 mo. MRIs at 12-14 years.



Behaviour?

Parallel studies have shown reduced IQs and cognitive and socioemotional deficits that are related to age at adoption and length of time in the SAME foster home.

Factors influencing brain development

- 1. sensory & motor experience
- 2. language & cognitive experience
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- **10. perinatal injury**
- 11. immune system

Play is a Form of Problembased Learning





What does play do to the brain?

- 1. increases the 'pruning' in prefrontal cortex.
- 2. Increases later flexibility (plasticity) in prefrontal cortex.

Bell et al., *Beh Brain Res*, 2010, 207, 7-13. Himmler et al., *Neurosc Lett*, 2013, 556, 42-45. Himmler et al., Synapse, in press.

Factors influencing brain development

- 1. sensory & motor experience
- 2. language & cognitive experience
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- **10. perinatal injury**
- **11. immune system**

Leung et al. BMC Pregnancy and Childbirth 2011, 11:12 http://www.biomedcentral.com/1471-2393/11/12

RESEARCH ARTICLE

Does prenatal micronutrient supplementation improve children's mental development? A systematic review

Brenda MY Leung $^{1,2^{\star}\dagger},$ Kristin P Wiens $^{2\dagger},$ Bonnie J Kaplan 1,2†

YES

BMC Pregnancy & Childbirth

Open Access

Factors influencing brain development

- 1. sensory & motor experience
- 2. language & cognitive experience
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- 10. perinatal injury
- **11. immune system**



The bugs influence the brain

Microorganisms can alter the Brain

Review

Nature Reviews Neuroscience 13, 701-712 (October 2012) | doi:10.1038,

Mind-altering microorganisms: the impact of the gut microbiota on brain and behaviour

John F. Cryan & Timothy G. Dinan

Biol Psychiatry, 2013 Nov 15;74(10):720-6. doi: 10.1016/j.biopsych.2013.05 Psychobiotics: a novel class of psychotropic. Dinan TG¹, Stanton C, Cryan JF.


Factors influencing brain development

- 1. sensory & motor experience
- 2. language & cognitive experience
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- 10. perinatal injury
- **11. immune system**

Perinatal Brain Injuries

Outcomes vary with age at injury, age at assessment, behaviour(s) measured, and sex.

The brain is always smaller but at some times there is compensation allowing the brain to do 'more with less.'



Perinatal Brain Injuries

Bottom line is that outcome is related to the precise time of brain development at injury (prenatal, infant, juvenile, early vs late adolescent, adult) and age at assessment.


There appear to be several distinct epochs... This is also true of cats, monkeys, & humans

See review by Kolb et al., *Progress in Brain Res*, 2013, 207, 35-64.

Perinatal Brain Injuries

Interventions make a difference: e.g., Tactile stimulation & complex housing.

See review by Kolb et al., *Progress in Brain Res*, 2013, 207, 35-64.





Housing pregnant moms in 'Condos' enhances Recovery from neonatal <u>mPFC</u> injury.



Perinatal Brain Injuries

Recovery is correlated with spontaneous neurogenesis, exuberant connections, synaptic space, & thalamic size.


See review by Kolb et al., *Progress in Brain Res*, 2013, 207, 35-64.

Factors influencing brain development

- 1. sensory & motor experience
- 2. language & cognitive experience
- 3. poverty
- 4. pre- and postnatal stress
- 5. psychoactive drugs
- 6. parent-child relationships
- 7. peer relationships
- 8. diet
- 9. gut bacteria
- **10.** perinatal injury
- **11. immune system**

Immune Proteins & Development

Many proteins 1st found in the immune system are also expressed in the developing CNS.




Activation of the immune system proposed in many disorders including ASD, Zika virus and schizophrenia.

e.g., Estes & McAllister, Science, 2016, 351, 919-920.

Cell Stem Cell Brief Report

Zika Virus Infects Human Cortical Neural Progenitors and Attenuates Their Growth

Hengli Tang, ^{1,1}, * Christy Hammack, ^{1,1}, Sarah C. Ogden, ^{1,11} Zhexing Wen, ^{2,3,11} Xuyu Qian, ^{2,4,11} Yujing Li,⁸ Bing Yao,⁸ Jaehoon Shin,^{2,4} Poiran Zhang,⁹ Emily M. Lee, ¹ Kimberly M. Christian,^{2,2} Ruth A. Didler,¹⁰ Peng Jin,⁹ Hongjun Song, ^{1,3,4,4,7} and (uc)-1 Ming^{2,3,4,7,4}



HEAD SIZE IN MICROCEPHALY







Baby with Typical Head Size

Baby with Microcephaly Seve

Conclusions

Brain development is complex & prolonged.

Brain plasticity in development is influenced by a wide range of early experiences including both pre- and postnatal experiences that may have good, bad, or ugly consequences.

